



☐ NOBLE NEW YORK ☐ NOBLE MISSISSIPPI Tel: 888-843-2040 Fax: 888-842-3977

Tel: 866-420-4041

Fax: 601-420-4040

Rheumatology

Delivery Nee	ed By: D	eliver to: 🗆 Patient	t's Home 🗆 Physici	an's Office 🗆 Othe	er		
PA	TIENT INFORM	ATION	PRES	CRIBER INFORM	ATION		
Address: City: Phone Number: _ Email Address: _ Last Four of Soc	State:		Office Contact Name Address: City: Phone Number: DEA/NPA #: PRESCRIPT	State: Fax:	Zip:		
Diamaraia			. INFORMATION	turata da maria cale fa			
Diagnosis:			Has the patient been treated previously for this condition?				
				□ Yes □ No			
Height: ft inches Weight: lbs							
		Date:	Other Notes:				
Allergies:							
			ON INFORMATION				
Medication:		e/Strength:	□ Inject SC every OTHER v	etions:	Quantity:	Refills:	
Actemra	□ 162mg/0.9ml Prefilled syringe □ 80 mg/4ml Vial □ 200mg/10ml Vial		□ Inject SC every OTHER V □ Inject SC every week Loading Dose: □ Infuse 4mg/kg	Maintenance Dose:	□ 4 week supply □ Other		
	□ 400mg/20ml Vial		(mg dose) via IV every 4 weeks	(mg dose) via IV every 4 weeks.			
Cimzia	□ 200 mg/ml Prefilled S' □ Starter Kit	YR	Loading Dose: □ Inject 400 mg SC at weeks 0, 2 and 4	Maintenance Dose: □ Inject 200 mg SC every other week □ Inject 400 mg SC every 4 weeks	☐ 4 week supply ☐ Other		
Cosentyx *Enhanced Specialty Pharmacy Program Participant	□ 150 mg Pen □ 150 mg SYR		Loading Dose: ☐ Inject 150 mg at weeks 0, 1, 2, 3, 4 ☐ Inject 400 mg at weeks 0, 1, 2, 3, 4	Maintenance Dose: □ Inject 150 mg every 4 weeks □ Inject 300 mg every 4 weeks	☐ 5 week supply (loading) ☐ 4 week supply (loading) ☐ Other		
Cosentyx *Enhanced Specialty Pharmacy Program Participant Covered Until You're Covered	□ 150 mg Pen □ 150 mg SYR		Loading Dose: □ Inject 150 mg at weeks 0, 1, 2, 3, 4 □ Inject 400 mg at weeks 0, 1, 2, 3, 4	Maintenance Dose: □ Inject 150 mg every 4 weeks □ Inject 300 mg every 4 weeks	☐ 5 week supply (loading) ☐ 4 week supply (loading) ☐ Other		
Enbrel Enbrel Mini Available	Standard: □ 25mg/0.5ml Prefilled SYR □ 50mg/ml Single Use Prefilled SYR □ 50mg/ml SureClick Autoinjector □ 25mg Vial	Mini: □ 50mg Enbrel Mini single dose prefilled cartridge	□ Inject 50mg SC TWICE a week (72-96 hours apart) □ Inject 50mg SC ONCE a week □ Inject 25mg SC TWICE a week (72-96 hours apart) □ Other		☐ 4 week supply☐ Other		
Humira	Standard: □ 40 mg/0.8ml Pen □ 40 mg/0.8ml Prefilled Syringe	Citrate-Free: □ 40mg/0.4 ml Pen □ 40 mg/0.4 ml Prefilled SYR	☐ Inject 40 mg SC every of ☐ Inject 40 mg SC once a v ☐ Other	☐ 4 week supply☐ Other☐			
☐ Patient is interested in patient support programs			□ Ancil	ary supplies provided for admi	nistration		
Phvs	ician Signature:			Date:			





☐ NOBLE NEW YORK ☐ NOBLE MISSISSIPPI Tel: 888-843-2040 Fax: 888-842-3977

Tel: 866-420-4041

Fax: 601-420-4040

Rheumatology

Delivery Need By: Deliver to: □ Patient's Home □ Physician's Office □ Other							
PA	TIENT INFORMATION		PRE	SCRIBER INFORM	IATION		
Address: City: Phone Number: _ Email Address: _ Last Four of Soci	Male Male Fem State: Zip: Zip:	nale Of Of Of Of Of De	fice Contact Nar Idress: :y: one Number: EA/NPA #:	State: Fax:	Zip:		
INSCRAI			ORMATION	HON CARD I K	ONT & DA		
Diagnosis:				en treated previously fo	r this condition?		
ICD-10 Code:			□ Yes □ No				
Height: ft inches Weight: lbs Last PPD Test: □ Positive □ Negative Date: Allergies:			Medications Failed: Medications On: Other Notes:				
		IPTION	INFORMATIC	N			
Medication:	Dosage/Strength:			rections:	Quantity:	Refills:	
Inflectra	□ 100 mg vial	□ 5ı m w	ading Dose: mg/kg (Dose ng) IV at 0, 2 and 6 reeks, then every 8 reeks thereafter	Maintenance Dose: □ 5mg/kg (Dose mg) IV every 8 weeks	□# of vials		
Kevzara	Prefilled Syringe: □ 150mg/1.14ml □ 200mg/1.14ml	□ 15	Prefilled Pen: ☐ 150mg/1.14ml ☐ 200mg/1.14ml		□ Inject mg once every TWO weeks	□ 4 week supply □ Other	
Olumiant	□ 2 mg tablet	□Та	ake one tablet (2mg)	☐ 4 week supply☐ Other	O tillo!		
Orencia	 □ 250mg Vial □ 125mg/ml SYR □ 125mg/ml Clickject □ 50 mg SYRINGE (for children > 2years and weighing 10kg to less than 25kg 	□ In w ev th	Dosing: Ifuse mg at reeks 0, 2, 4 and reey 4 weeks reereafter	Subcutaneous Dosing: □ Inject 125 mg subcutaneously once a week	□ 4 week supply □ Other		
Otezla	□ Starter Kit		ther rter Kit:	Maintenance Dose:			
	□ 30 mg	□ Ta	ake as directed	□ take 30 mg twice daily			
Otrexup	□ 10 mg / 0.4 ml autoinjector □ 12.5 mg/ 0.4 ml autoinjector □ 15mg/ 0.4 ml autoinjector □ 17.5mg/ 0.4 ml autoinjector □ 20mg/ 0.4 ml autoinjector □ 22.5mg/ 0.4 ml autoinjector □ 25mg/ 0.4 ml autoinjector		□ Inject mg subcutaneously once weekly □ Other		☐ 4 week supply☐ Other		
Rasuvo	□ 7.5 mg/0.15ml autoinjector □ 10 mg/0.2ml autoinjector □ 12.5 mg/0.25ml autoinjector □ 15 mg/.3ml autoinjector □ 17.5 mg/0.35ml autoinjector □ 20 mg/0.4ml autoinjector □ 22.5 mg/0.45ml autoinjector □ 25 mg/0.5ml autoinjector □ 30 mg/0.6ml autoinjector		ject mg suk ther	☐ 4 week supply☐ Other			
Rayos	□ 1 mg tablet □ 2 mg tablet □ 5 mg tablet		☐ Take mg by mouth once per day ☐ Other ☐ Applies provided for admit		☐ 4 week supply☐ Other		
	it is interested in patient support programs		□ Ar	ncillary supplies provided for adm	IIIISTRATION		
Phvs	ician Signature:			Date:			





☐ NOBLE NEW YORK ☐ NOBLE MISSISSIPPI Tel: 888-843-2040 Fax: 888-842-3977

Tel: 866-420-4041

Fax: 601-420-4040

Rheumatology

Delivery Nee	ed By: De	liver to: 🗆 Patient	's Home 🗆 Physicia	an's Office 🗆 Othe	er	
PATIENT INFORMATION PRESCRIBER INFORMATION						
Patient Name:			Prescriber's Name:			
INSURAN	NCE - PLEASE	FAX COPY O	F PRESCRIPTI		ONT & BA	CK
		CLINICAL	INFORMATION			
Diagnosis: ICD-10 Code: Height: ft inches Weight: lbs Last PPD Test: □ Positive □ Negative Date:			Has the patient been treated previously for this condition? Yes No Medications Failed: Medications On: Other Notes:			
		PRESCRIPTION	DN INFORMATION			
Medication:	Dosage/	Strength:	,	tions:	Quantity:	Refills:
Remicade	□ 100mg vial		Loading Dose: □ Infuse 5mg/kg (Dose mg) via IV at 0, 2 and 6 weeks □ Infuse mg every _ □ Other	IVevery 8 weeks		
Renflexis	□ 100mg vial		Loading Dose: ☐ Infuse 5mg/kg (Dosemg) via IV at 0, 2 and 6 weeks, then every 8 weeks thereafter	Maintenance Dose: □ Infuse 5mg/kg (Dose mg) IV every 8 weeks	U# of Vials	
Rituxan	☐ 100 mg/10ml vial ☐ 500 mg /50ml vial		□ IV mg every weeks □ Specified:		uials # of	
Simpon®	Prefilled Syringe: ☐ 50ml/0.5ml ☐ 100mg/1ml	SmartJect AutoInjector: □ 50ml/0.5ml □ 100mg/1ml	□ Inject 100 mg subcutaneously once a month □ Inject 50 mg subcutaneously once a month		☐ 4 week supply☐ Other	
Simponi Aria	□ 50mg/4ml single dose vial		Loading Dose: ☐ Infuse 2mg/kg (Dose mg) via IV over 30 minutes at weeks 0 and 4	Maintenance Dose: □ Infuse 2mg/kg (Dose mg) via IV over 30 minutes every 8 weeks		
Taltz	□ 80 ml/ml single-dose prefilled autoinjector □ 80mg/ml single-dose prefilled syringe		Loading Dose: □ Inject 160mg subcutaneously at week zero	Maintenance Dose: □ Inject 80mg subcutaneously every 4 weeks	☐ pens ☐ syringes ☐ Other	
Xeljanz	□ 5 mg tablet		□ Take one tablet twice a day		☐ 4 week supply☐ Other	
Xeljanz XR	□ 11 mg tablet		☐ Take one tablet once a day		☐ 4 week supply☐ Other	
Other						
□ Patier	nt is interested in patient supp	ort programs	□ Ancill	ary supplies provided for adm	inistration	
Phys	sician Signature:			Date:		